

# Derivatives and functional linear models

A first look at a differential equation  
as a modelling tool.

*The oil refinery data*

*The melanoma data*

*Summary*

*Home Page*

*Title Page*

◀ ▶

◀ ▶

Page 1 of 18

*Go Back*

*Full Screen*

*Close*

*Quit*

# 1. The oil refinery data

*The oil refinery data*

*The melanoma data*

*Summary*

*Home Page*

*Title Page*

◀◀ ▶▶

◀ ▶

Page 2 of 18


*Go Back*

*Full Screen*

*Close*

*Quit*

# Refinery output $x(t)$ (top panel) and input $u(t)$ (bottom panel)



The oil refinery data

The melanoma data

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 3 of 18

Go Back

Full Screen

Close

Quit

- This is a simple input/output system in an oil refinery in Corpus Christi, Texas.
- A fluid, called *reflux*, flows into *tray 47* in a distillation column in an oil refinery.
- The input variable  $u(t)$  is the flow rate.
- The fluid level in the tray is the output variable  $x(t)$ .

The oil refinery data

The melanoma data

Summary

Home Page

Title Page

◀◀ ▶▶

◀ ▶

Page 4 of 18

Go Back

Full Screen

Close

Quit

# Variation on two time scales

- Over the longer scale, tray level changes from 0 to around 4.
- But we are also interested in how rapidly the change takes place; that is, short-scale variation.
- It looks like about  $\frac{2}{3}$  of the change takes about 50 minutes, and the final value is reached in 200 minutes or so.

*The oil refinery data*

*The melanoma data*

*Summary*

*Home Page*

*Title Page*

◀ ▶

◀ ▶

Page 5 of 18

*Go Back*

*Full Screen*

*Close*

*Quit*

# A concurrent functional regression

- We can model the output *state*  $\text{Tray}(t)$  as a simple time-varying regression on input state  $\text{Reflux}(t)$ :

$$\text{Tray}(t) = \text{Reflux}(t)\beta(t) + \epsilon(t)$$

- In functional data analysis, we call this a *concurrent* regression because only the simultaneous influence of the input on the output is modelled.
- A least squares estimate  $\hat{\beta}(t)$  of the regression coefficient function minimizes

$$\text{SSE} = \int [\text{Tray}(t) - \text{Reflux}(t)\beta(t)]^2 dt$$

- $\hat{\beta}(t)$  is modelled with an expansion in terms of several B-spline basis functions.

The oil refinery data

The melanoma data

Summary

Home Page

Title Page

◀ ▶

◀ ▶


Page 6 of 18

Go Back

Full Screen

Close

Quit



But does this tell us anything new? The regression function is just as complicated as the output.

[The oil refinery data](#)

[The melanoma data](#)

[Summary](#)

[Home Page](#)

[Title Page](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

# Making the derivative $D\text{Tray}(t)$ the output

- We now model the *rate of change*  $D\text{Tray}(t)$ , using the output state  $\text{Tray}(t)$  and the input  $\text{Reflux}(t)$  as covariates.
- We'll use constants for the two regression functions:

$$D\text{Tray}(t) = -\beta_1\text{Tray}(t) + \beta_2\text{Reflux}(t) + \epsilon(t)$$

- This is an example of a *first order differential equation with constant coefficients*.
- We see that it is just another form of concurrent functional linear model, now with two covariates, but with regression functions  $\beta_1$  and  $\beta_2$  that are constant.

The oil refinery data

The melanoma data

Summary

Home Page

Title Page

◀ ▶

◀ ▶

Page 8 of 18


Go Back

Full Screen

Close

Quit





The solid line is the derivative estimated from the data, and the dashed line is the model's fit to this derivative.  $\hat{\beta}_1 = 0.02$  and  $\hat{\beta}_2 = 0.19$

[The oil refinery data](#)

[The melanoma data](#)

[Summary](#)

[Home Page](#)

[Title Page](#)

[«](#) [»](#)

[◀](#) [▶](#)

Page 9 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

- A differential equation that is this simple has an explicit solution.

$$x(t) = e^{-\beta_1 t} [x(0) - (\beta_2/\beta_1) \int_0^t e^{\beta_1 s} u(s) ds].$$

- $\beta_1 \approx 0.02$  is the rate constant, and therefore controls the rate of change of  $\text{Tray}$  level. About 2/3 of the change takes  $1/\beta_1$  time units, and the final level is nearly reached in  $4/\beta_1$  time units.  $\beta_1$  models the *dynamic behavior* of  $\text{Tray}$ .
- $\beta_2 \approx 0.19$ , along with  $\beta_1$ , defines the ultimate change; the long-term *gain* per unit increase in Reflux flow is  $\beta_2/\beta_1 \approx 9.5$ .

The oil refinery data

The melanoma data

Summary

Home Page

Title Page

◀ ▶

◀ ▶


Page 10 of 18

Go Back

Full Screen

Close

Quit



Modelling the rate of change  $D_{\text{Tray}}(t)$  directly produces a fine fit to the data with only two parameters.

## 2. The melanoma data

*The oil refinery data*

*The melanoma data*

*Summary*

*Home Page*

*Title Page*

◀◀ ▶▶

◀ ▶

Page 12 of 18

*Go Back*


*Full Screen*

*Close*

*Quit*

# Age-adjusted melanoma incidences for Connecticut.

The solid line is a spline smooth with penalty on  $D^4x$ .



[The oil refinery data](#)

[The melanoma data](#)

[Summary](#)

[Home Page](#)

[Title Page](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

Page 13 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

# Estimating a differential operator $L$

- A differential operator  $L$  is just a re-arranged differential equation.
- Can we smooth the data and estimate the operator at the same time?
- Will this give us a better fit to the data with fewer degrees of freedom used up?
- We will try the operator

$$Lx = \beta_1 D^2 x + D^4 x$$

- This operator a tilted line plus sinusoidal trend with the period to be estimated from the data.

# The algorithm

- Start with  $\beta_1 = 0$  and  $Lx = D^4x$ , and estimate derivatives up to order 4, choosing  $\lambda$  to minimize the GCV criterion.
- Carry out concurrent functional regression to estimate  $\beta_1$ .
- Re-smooth using  $Lx$ , again re-computing derivatives and minimizing GCV.
- Continue until the parameter estimates converge.

*The oil refinery data*

*The melanoma data*

*Summary*

*Home Page*

*Title Page*

*◀ ▶*

*◀ ▶*

*Page 15 of 18*


*Go Back*

*Full Screen*

*Close*


*Quit*

Solid blue line is smooth using converged  $Lx$  penalty. Dashed line is smoothing using  $D^4$  penalty. Dotted line is a solution to differential equation  $Lx = 0$ .

[The oil refinery data](#)[The melanoma data](#)[Summary](#)[Home Page](#)[Title Page](#)[◀](#) [▶](#)[◀](#) [▶](#)[Page 16 of 18](#)[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)



# Phase-plane plot



[The oil refinery data](#)

[The melanoma data](#)

[Summary](#)

[Home Page](#)

[Title Page](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

Page 17 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

### 3. Summary

- By using a derivative as the dependent variable, we can model the rate of change of an output variable.
- The resulting differential equation can be solved to provide a model for the output variable itself.
- We effectively get two or more models for the price of one.
- We are simultaneously modelling the *level* or *state* of a system and its *dynamics*.

[Home Page](#)[Title Page](#)[◀◀](#) [▶▶](#)[◀](#) [▶](#)

Page 18 of 18

[Go Back](#)[Full Screen](#)[Close](#)[Quit](#)